

Spin analog of the controlled Josephson charge current

Sudhansu S. Mandal and Soumya P. Mukherjee

Theoretical Physics Department, Indian Association for the Cultivation of Science, Jadavpur, Kolkata 700 032, India

(Dated: February 4, 2008)

We propose a controlled Josephson spin current across the junction of two non-centrosymmetric superconductors like CePt₃Si. The Josephson spin current arises due to direction dependent tunneling matrix element and different momentum dependent phases of the triplet components of the gap function. Its modulation with the angle ξ between the noncentrosymmetric axes of two superconductors is proportional to $\sin \xi$. This particular dependence on ξ may find application of the proposed set-up in making a *Josephson spin switch*.

PACS numbers: 74.50.+r, 72.25.-b, 74.20.Rp, 74.70.Tx

Traditionally Josephson junctions [1, 2] in superconductors draw interest both scientifically and its applicability in making devices. With no exception, they have also been studied in unconventional superconductors like spin-singlet cuprate [3] and spin-triplet Sr₂RuO₄ [4] superconductors. However no Josephson junction between nonmagnetic superconductors is known to generate spin-polarized current. The purpose of this letter is to theoretically show that the direction dependent tunneling matrix element across the junction between two recently discovered [5] non-centrosymmetric superconductors like CePt₃Si, leads to tunneling of both spin-singlet and spin-triplet Cooper pairs. As a consequence, nonvanishing spin-Josephson current is viable along with the usual charge-Josephson current. This *novel* spin-Josephson current depends on the relative angle ξ between the axes of non-centrosymmetry $\hat{n}_L$ and $\hat{n}_R$ in the left and right side of the junction respectively. This angular dependence may be used to make a *Josephson spin switch*.

The normal state Hamiltonian [6, 7] for the electrons in a band of a lattice without inversion symmetry is

$$H_0 = \sum_{\mathbf{k}, s} \xi_{\mathbf{k}} c_{\mathbf{k}s}^\dagger c_{\mathbf{k}s} + \sum_{\mathbf{k}, s, s'} \mathbf{g}_{\mathbf{k}} \cdot \boldsymbol{\sigma}_{ss'} c_{\mathbf{k}s}^\dagger c_{\mathbf{k}s'}, \quad (1)$$

where electrons with momentum $\mathbf{k}$ and spin s ($\uparrow$ or $\downarrow$) are created (annihilated) by the operators $c_{\mathbf{k}s}^\dagger$ ($c_{\mathbf{k}s}$), and $\xi_{\mathbf{k}}$ is the band energy measured from the Fermi energy ϵ_F . The second term in the Hamiltonian (1) breaks parity symmetry as $\mathbf{g}_{-\mathbf{k}} = -\mathbf{g}_{\mathbf{k}}$ for a non-centrosymmetric system. For a system like Heavy fermion compound CePt₃Si which has layered structure, H_0 is considered to be two-dimensional. For such a system of electrons with band mass m , $\xi_{\mathbf{k}} = \frac{k^2}{2m} - \epsilon_F$ and $\mathbf{g}_{\mathbf{k}} = \alpha \boldsymbol{\eta}_{\mathbf{k}}$ where $\boldsymbol{\eta}_{\mathbf{k}} = \hat{n} \times \mathbf{k}$, i.e., the spin-orbit interaction is of Rashba type [8] and α is the Rashba parameter. Here $\hat{n}$ represents the axis of non-centrosymmetry which is perpendicular to the plane of the system. Due to the breaking down of the parity, spin degeneracy of the band is lifted; by diagonalizing H_0 , one finds two spin-split bands with energies $\xi_{\mathbf{k}\lambda} = \xi_{\mathbf{k}} + \lambda \alpha |\mathbf{k}|$ where $\lambda = \pm$ describes helicity of the spin-split bands. Therefore in the diagonalized basis H_0 (1) becomes $H_0 = \sum_{\mathbf{k}, \lambda=\pm} \xi_{\mathbf{k}\lambda} \tilde{c}_{\mathbf{k}\lambda}^\dagger \tilde{c}_{\mathbf{k}\lambda}$, where

$\tilde{c}_{\mathbf{k}\lambda} = (c_{\mathbf{k}\uparrow} - i\lambda \exp(i\phi_{\mathbf{k}}) c_{\mathbf{k}\downarrow}) / \sqrt{2}$ is the electron destruction operator and $\tilde{c}_{\mathbf{k}\lambda}^\dagger = (c_{\mathbf{k}\uparrow}^\dagger + i\lambda \exp(-i\phi_{\mathbf{k}}) c_{\mathbf{k}\downarrow}^\dagger) / \sqrt{2}$ is the electron creation operator in band λ with momentum $\mathbf{k}$ whose orientation with $\hat{x}$ -axis is $\phi_{\mathbf{k}}$. The density of electronic states at Fermi energy in these bands may be found as $\nu_{\lambda} = \frac{m}{2\pi} \left(1 - \lambda m \alpha / \sqrt{k_F^2 + m^2 \alpha^2}\right)$, where $k_F = \sqrt{2m\epsilon_F}$ is the Fermi momentum.

Band structure calculation [9] on CePt₃Si reveals that the energy difference between two spin-split bands near k_F is 50–200 meV which is much larger than the superconducting critical temperature, $k_B T_c \approx 0.06$ meV [5]. The formation of Cooper pairing between electrons in different spin-split bands may thus be ignored. The Hamiltonian for these superconductors may then be written as

$$H_1 = \sum_{\mathbf{k}, \lambda=\pm} \left[\xi_{\mathbf{k}\lambda} \tilde{c}_{\mathbf{k}\lambda}^\dagger \tilde{c}_{\mathbf{k}\lambda} + \left(\Delta_{\mathbf{k}\lambda} \tilde{c}_{\mathbf{k}\lambda}^\dagger \tilde{c}_{-\mathbf{k}\lambda}^\dagger + \text{h.c.} \right) \right], \quad (2)$$

where Cooper pairs are only between intraband electrons. Therefore the normal and anomalous Green's functions are obtained respectively as $\mathcal{G}_{\lambda}(\mathbf{k}, i\epsilon_n) = -(i\epsilon_n + \xi_{\mathbf{k}\lambda}) / (\epsilon_n^2 + \xi_{\mathbf{k}\lambda}^2 + |\Delta_{\mathbf{k}\lambda}|^2)$ and $\mathcal{F}_{\lambda}(\mathbf{k}, i\epsilon_n) = \Delta_{\mathbf{k}\lambda} / (\epsilon_n^2 + \xi_{\mathbf{k}\lambda}^2 + |\Delta_{\mathbf{k}\lambda}|^2)$, where ϵ_n is the fermionic Matsubara frequency [13]. The superconducting order parameter $\Delta_{\mathbf{k}\lambda}$ obeys the symmetry [9, 10]: $\Delta_{-\mathbf{k}\lambda} = -\Delta_{\mathbf{k}\lambda}$. We consider $\Delta_{\mathbf{k}\lambda} = \tilde{\Delta}_{\mathbf{k}\lambda} \Lambda_{\mathbf{k}}$, i.e., the angular dependence of $\mathbf{k}$ on the order parameters of two bands are assumed to be same. Apart from the overall phase rigidity angle Θ of the superconductor, there may be a relative phase difference θ between superconductors of two bands: $\tilde{\Delta}_{\mathbf{k}+} = |\tilde{\Delta}_{\mathbf{k}+}| e^{i\Theta}$ and $\tilde{\Delta}_{\mathbf{k}-} = |\tilde{\Delta}_{\mathbf{k}-}| e^{i(\Theta+\theta)}$. Reverting H_1 (2) to spin-up (-down) basis, we find

$$H_1 = \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^\dagger \begin{pmatrix} \xi_{\mathbf{k}} & \Delta_{\mathbf{k},\uparrow\uparrow} & \Gamma_{\mathbf{k},R} & \Delta_{\mathbf{k},\uparrow\downarrow} \\ \Delta_{\mathbf{k},\uparrow\uparrow}^* & -\xi_{\mathbf{k}} & \Delta_{\mathbf{k},\uparrow\downarrow}^* & \Gamma_{\mathbf{k},R} \\ \Gamma_{\mathbf{k},R}^* & \Delta_{\mathbf{k},\uparrow\downarrow} & \xi_{\mathbf{k}} & \Delta_{\mathbf{k},\downarrow\downarrow} \\ \Delta_{\mathbf{k},\uparrow\downarrow}^* & \Gamma_{\mathbf{k},R} & \Delta_{\mathbf{k},\downarrow\downarrow}^* & -\xi_{\mathbf{k}} \end{pmatrix} \Psi_{\mathbf{k}}, \quad (3)$$

where $\Delta_{\mathbf{k},\uparrow\uparrow} = \frac{1}{2}(\tilde{\Delta}_{\mathbf{k}+} + \tilde{\Delta}_{\mathbf{k}-})\Lambda_{\mathbf{k}}$, $\Delta_{\mathbf{k},\downarrow\downarrow} = \frac{1}{2} \exp[2i\phi_{\mathbf{k}}](\tilde{\Delta}_{\mathbf{k}+} + \tilde{\Delta}_{\mathbf{k}-})\Lambda_{\mathbf{k}}$, and $\Delta_{\mathbf{k},\uparrow\downarrow} = -\Delta_{\mathbf{k},\downarrow\uparrow} = \frac{i}{2} \exp[i\phi_{\mathbf{k}}](\tilde{\Delta}_{\mathbf{k}+} - \tilde{\Delta}_{\mathbf{k}-})\Lambda_{\mathbf{k}}$ are the different components of pairing potential $\Delta_{\mathbf{k},ss'}$ between electrons with




FIG. 1: Schematic picture of two non-centrosymmetric superconductors, denoted by L and R , coupled at $x = x_0$. Their orientations are characterized by the axes of non-centrosymmetry, $\hat{n}_L$ and $\hat{n}_R$.

spins s and s' , and $\Gamma_{\mathbf{k},R} = i\alpha|\mathbf{k}|\exp[-i\phi_{\mathbf{k}}]$ is the Rashba spin-orbit coupling potential. The Hamiltonian (3) has been expressed in the basis such that $\Psi_{\mathbf{k}}^\dagger = (c_{\mathbf{k}\uparrow}^\dagger, c_{-\mathbf{k}\uparrow}^\dagger, c_{\mathbf{k}\downarrow}^\dagger, c_{-\mathbf{k}\downarrow}^\dagger)$. Since $\Delta_{\mathbf{k},\uparrow\downarrow} = -\Delta_{-\mathbf{k},\uparrow\downarrow}$, there is no triplet component with zero projection along the spin-quantization direction; pairing between electrons with unequal spins entirely gives rise to singlet component. If we choose $\Lambda_{\mathbf{k}} = -i\exp[-i\phi_{\mathbf{k}}]$, the triplet component of the pairing may be expressed as $\hat{\Delta}_{\mathbf{k},T} = (\mathbf{d}_{\mathbf{k}} \cdot \boldsymbol{\sigma})i\sigma_y$ with $\mathbf{d}_{\mathbf{k}} = \frac{1}{2|\mathbf{k}|}(\tilde{\Delta}_{k+} + \tilde{\Delta}_{k-})\boldsymbol{\eta}_{\mathbf{k}}$. In other words, the only stable [7] spin triplet component whose $\mathbf{d}_{\mathbf{k}} \parallel \boldsymbol{\eta}_{\mathbf{k}}$ corresponds to this $\Lambda_{\mathbf{k}}$. Therefore $\Delta_{\mathbf{k},\uparrow\uparrow}$ and $\Delta_{\mathbf{k},\downarrow\downarrow}$ will have equal but opposite momentum-dependent phase. Accordingly the singlet component of the pairing potential becomes $\Delta_{\mathbf{k},S} = \frac{1}{2}(\tilde{\Delta}_{k+} - \tilde{\Delta}_{k-})$. If $|\tilde{\Delta}_{k+}| \neq |\tilde{\Delta}_{k-}|$, the admixture [6, 7] of singlet and triplet pairing takes pace. Recent observation [11] of Josephson current in the junction of CePt₃Si and s -wave superconductor suggests the existence of spin-singlet order parameter, while larger [5] upper critical field H_{c2} seems to suggest that the spin-triplet pairing occurs. One normally assumes the superconducting order parameter to be independent of the magnitude of momentum, i.e., $\tilde{\Delta}_{k\lambda}$ is k -independent. The anomalous Green's function $\mathcal{F}_{ss'}(\mathbf{k}, i\epsilon_n)$ in the up-down basis is related to $\mathcal{F}_\lambda$ as $\mathcal{F}_{\uparrow\uparrow} = \frac{1}{2}(\mathcal{F}_+ + \mathcal{F}_-)$, $\mathcal{F}_{\downarrow\downarrow} = \frac{1}{2}e^{2i\phi_{\mathbf{k}}}(\mathcal{F}_+ + \mathcal{F}_-)$, and $\mathcal{F}_{\uparrow\downarrow} = -\mathcal{F}_{\downarrow\uparrow} = \frac{i}{2}e^{i\phi_{\mathbf{k}}}(\mathcal{F}_+ - \mathcal{F}_-)$. Apart from the applicable momentum-dependent phases, triplet (singlet) component of $\mathcal{F}_{ss'}$ is the addition (subtraction) of anomalous Green's function of two spin-split bands.

We now consider Josephson tunneling between two such non-centrosymmetric superconductors as depicted in Fig. 1. The Hamiltonian for the system then reads $H = H_L + H_R + H_T$, where H_L and H_R are the bulk Hamiltonian of left and right side of the junction

respectively and H_T describes the tunneling between these two sides. The Hamiltonian H_L is described by H_1 (2) and H_R is also defined equivalently with the change in the notation of momentum $\mathbf{k} \rightarrow \mathbf{p}$ to distinguish each side of the junction. Further, each parameter in the left (right) is denoted by superscript or subscript L (R). The tunneling Hamiltonian reads $H_T = \sum_{\mathbf{k}\mathbf{p},s} (T_{\mathbf{k}\mathbf{p}}c_{\mathbf{k}s}^\dagger c_{\mathbf{p}s} + T_{\mathbf{k}\mathbf{p}}^*c_{\mathbf{p}s}^\dagger c_{\mathbf{k}s})$, where $T_{\mathbf{k}\mathbf{p}}$ is the tunneling matrix element for an electron with momentum $\mathbf{p}$ to tunnel from right side to left side with momentum $\mathbf{k}$. Time-reversal symmetry of H_T suggests $T_{-\mathbf{k},-\mathbf{p}} = T_{\mathbf{k}\mathbf{p}}^*$.

In a charge-tunneling process, number of electrons in either side of the junction changes with time, e.g., the rate of change in the number of electrons of the right side of the junction $N_R = \sum_{\mathbf{p},s} c_{\mathbf{p}s}^\dagger c_{\mathbf{p}s}$ reads as $\dot{N}_R = i[H_T, N_R]$. Therefore the tunneling charge-current becomes $I_c(t) = -e\langle \dot{N}_R(t) \rangle$: one part corresponds to quasi-particle tunneling and the other part is due to process of Josephson tunneling [12, 13], i.e., tunneling of the Cooper pairs. We are herewith interested in the Josephson charge current defined as $I_J^c(t) = 2e\Im[\exp(-2ieVt)\Phi_{\text{ret}}^c(eV)]$ at time t and for an applied bias voltage V across the junction, where $\Phi^c(i\omega_m) = -\int_0^\beta d\tau e^{i\omega_m\tau} \langle T_\tau A(\tau)A(0) \rangle$ with imaginary time τ , $A(\tau) = \sum_{\mathbf{k}\mathbf{p},s} T_{\mathbf{k}\mathbf{p}}c_{\mathbf{k}s}^\dagger(\tau)c_{\mathbf{p}s}(\tau')$, the bosonic Matsubara frequency [13] ω_m , and inverse temperature β . Instead of the axes of noncentrosymmetry of two sides of the junction being parallel, general axes $\hat{n}_L$ and $\hat{n}_R$ will lead to $\hat{\boldsymbol{\eta}}_{\mathbf{k}} = \hat{n}_L \times \hat{\mathbf{k}}$ and $\hat{\boldsymbol{\eta}}_{\mathbf{p}} = \hat{n}_R \times \hat{\mathbf{p}}$. In that case, Josephson tunneling occurs in both singlet and triplet channels. Separating these channels, we find

$$\Phi^c(i\omega_m) = - \sum_{i\epsilon_n, \mathbf{k}\mathbf{p}} \frac{|T_{\mathbf{k}\mathbf{p}}|^2}{2\beta} [\chi_S + (\hat{\boldsymbol{\eta}}_{\mathbf{k}} \cdot \hat{\boldsymbol{\eta}}_{\mathbf{p}})\chi_T], \quad (4)$$

where

$$\chi_S = \sum_{\lambda, \lambda'} \lambda \lambda' \Lambda_{\mathbf{k}} \Lambda_{\mathbf{p}}^* \mathcal{F}_\lambda^\dagger(\mathbf{k}, i\epsilon_n) \mathcal{F}_{\lambda'}(\mathbf{p}, i\epsilon_n - i\omega_m) \quad (5)$$

and

$$\chi_T = \sum_{\lambda, \lambda'} \Lambda_{\mathbf{k}} \Lambda_{\mathbf{p}}^* \mathcal{F}_\lambda^\dagger(\mathbf{k}, i\epsilon_n) \mathcal{F}_{\lambda'}(\mathbf{p}, i\epsilon_n - i\omega_m) \quad (6)$$

represent singlet and triplet contributions respectively. While the direction independent electronic tunneling probability $|T_{\mathbf{k}\mathbf{p}}|^2 = |T|^2$ leads to tunneling of singlet component only, the triplet components of Cooper pairs tunnel due to realistic direction dependent [14] $|T_{\mathbf{k}\mathbf{p}}|^2$. This triplet part in the charge Josephson tunneling process may effectively provide spin Josephson tunneling. The rate of change of spin $S_z = \sum_{\mathbf{p}} [c_{\mathbf{p}\uparrow}^\dagger c_{\mathbf{p}\uparrow} - c_{\mathbf{p}\downarrow}^\dagger c_{\mathbf{p}\downarrow}]$ due to tunneling, i.e., $\dot{S}_z = i[H_T, S_z]$, gives rise to spin-Josephson tunneling current $I_J^s(t) = -2\Im[\exp(-2ieVt)\Phi_{\text{ret}}^s(eV)]$ where

$\Phi^s(i\omega_m) = -\int_0^\beta d\tau e^{i\omega_m\tau} \langle T_\tau B(\tau) A(0) \rangle$ with $B(\tau) = \sum_{\mathbf{k}\mathbf{p}} T_{\mathbf{k}\mathbf{p}} [c_{\mathbf{k}\uparrow}^\dagger(\tau) c_{\mathbf{p}\uparrow}(\tau) - c_{\mathbf{k}\downarrow}^\dagger(\tau) c_{\mathbf{p}\downarrow}(\tau)]$. We thus find

$$\Phi^s(i\omega_m) = i \sum_{i\epsilon_n, \mathbf{k}\mathbf{p}} \frac{|T_{\mathbf{k}\mathbf{p}}|^2}{2\beta} \sum_j (\hat{\eta}_k \times \hat{\eta}_p)_j \chi_T, \quad (7)$$

where $(\hat{\eta}_k \times \hat{\eta}_p)_j$ represents j -th spatial component of $\hat{\eta}_k \times \hat{\eta}_p$.

We assume that the non-centrosymmetric axes are parallel to the interface of the junction as shown in Fig. 1. The probability of tunneling [14] will be the most along the direction transverse to the interface. In terms of normal state electronic tunneling conductance $G_N = I_N/V$ with I_N being normal state tunneling current, we find dc ($V = 0$) charge and spin Josephson currents:

$$I_J^c = \left(\frac{G_N}{e\mathcal{F}} \right) [\sin \psi (g_+ \mathcal{A}_1 + g_- \mathcal{A}_2) + \cos \psi (g_+ \mathcal{A}_3 + g_- \mathcal{A}_4)] \quad (8)$$

$$I_J^s = \left(\frac{G_N}{e^2 \mathcal{F}} \right) \delta \kappa \sin \xi [\cos \psi (\mathcal{A}_1 + \mathcal{A}_2) - \sin \psi (\mathcal{A}_3 + \mathcal{A}_4)], \quad (9)$$

where $\psi = \Theta_L - \Theta_R$, $g_\pm = \delta \cos \xi \pm 1$, $\delta \in (0, 1)$ is a parameter depending on the model [15] of tunneling matrix element, i.e., the angular dependence of $T_{\mathbf{k}\mathbf{p}}$, $\cos \xi = \hat{n}_L \cdot \hat{n}_R$ where both $\hat{n}_L$ and $\hat{n}_R$ are in the plane of the interface, and κ is the projection of $\hat{n}_L \times \hat{n}_R$ along perpendicular to the interface. Further $\mathcal{F} = \pi \sum_{\lambda\lambda'} \nu_\lambda^L \nu_{\lambda'}^R (1 + \delta \lambda \lambda' \cos \xi)$ describes the dependence of normal state tunneling current on δ , $\mathcal{A}_1 = \Gamma_{++} + \Gamma_{--} \cos(\theta_L - \theta_R)$, $\mathcal{A}_2 = \Gamma_{-+} \cos \theta_L + \Gamma_{+-} \cos \theta_R$, $\mathcal{A}_3 = \Gamma_{--} \sin(\theta_L - \theta_R)$, and $\mathcal{A}_4 = \Gamma_{-+} \sin \theta_L - \Gamma_{+-} \sin \theta_R$ with

$$\Gamma_{\lambda\lambda'} = \pi \nu_\lambda^L \nu_{\lambda'}^R \frac{|\tilde{\Delta}_\lambda^L| |\tilde{\Delta}_{\lambda'}^R|}{|\tilde{\Delta}_\lambda^L| + |\tilde{\Delta}_{\lambda'}^R|} K \left(\frac{|\tilde{\Delta}_\lambda^L| - |\tilde{\Delta}_{\lambda'}^R|}{|\tilde{\Delta}_\lambda^L| + |\tilde{\Delta}_{\lambda'}^R|} \right) \quad (10)$$

at zero temperature, where K is the elliptic function of first kind.

Charge (8) and spin (9) dc Josephson currents depend on both sine and cosine of the global phase difference ψ between two superconductors. Instead of continuous relative phase difference between superconductors in each spin-split band, there is a possibility of phase locking such that $\theta_L, \theta_R = n\pi$ ($n \in \mathbb{Z}$). In this case, $\mathcal{A}_3 = \mathcal{A}_4 = 0$ and hence $I_J^c(V=0) \propto \sin \psi$ and $I_J^s(V=0) \propto \cos \psi$. This situation is also true for $\theta_L = \theta_R$, even in the absence of their locking at $n\pi$. The critical charge and spin Josephson currents are modulated with the angle ξ .

For general values of θ_L and θ_R , charge (8) and spin (9) Josephson currents may be parametrized as $I_J^c = J_1^c \sin(\psi + \chi_1) + J_2^c \sin(\psi + \chi_2)$ and $I_J^s = J^s \cos(\psi + \chi_1)$, where $J_1^c(e\mathcal{F}/G_N) = \mathcal{C} \delta \cos \xi$, $J_2^c(e\mathcal{F}/G_N) = \mathcal{D}$, $J^s(e^2 \mathcal{F}/G_N) = \mathcal{C} \delta \kappa \sin \xi$, $\chi_1 = \tan^{-1}(\frac{\mathcal{A}_3 + \mathcal{A}_4}{\mathcal{A}_1 + \mathcal{A}_2})$ and $\chi_2 = \tan^{-1}(\frac{\mathcal{A}_3 - \mathcal{A}_4}{\mathcal{A}_1 - \mathcal{A}_2})$ with $\mathcal{C} = [(\mathcal{A}_1 + \mathcal{A}_2)^2 + (\mathcal{A}_3 + \mathcal{A}_4)^2]^{1/2}$,

and $\mathcal{D} = [(\mathcal{A}_1 - \mathcal{A}_2)^2 + (\mathcal{A}_3 - \mathcal{A}_4)^2]^{1/2}$. With the application of external flux, current and phase relationship [2, 3] for I_J^c may be found out. This will determine J_1^c and J_2^c and thereby δ and ξ . These determinations will predict the value of I_J^s . The spin current may be tested if the proposed set-up acts as a source of spin current.

The spin-Josephson current (9) is proportional to $\sin \xi$ which means I_J^s is maximum when the orientation of the axes of non-centrosymmetry between left and right side of the superconductors are transverse, and it vanishes when the axes are parallel. This particular property should be useful to control spin-Josephson current by orienting the axes about which the inversion symmetries are lost. Therefore it will be useful to build a *Josephson spin switch*.

To summarize, we have shown that the Josephson tunneling process between two superconductors without inversion symmetry consists of tunneling of both spin-singlet and spin-triplet Cooper pairs. Because of the direction dependent tunneling matrix element in real junction and the momentum dependent phases of the triplet components of the gap function, *viz.* $\Delta_{\mathbf{k},\uparrow\uparrow}$ and $\Delta_{\mathbf{k},\downarrow\downarrow}$, are opposite, there is a net spin-Josephson current. This spin current is proportional to sine of the angle ξ between the axes of non-centrosymmetry in the bulk superconductors in each side of the junction. It may be used to make a *Josephson spin switch* with the controlling parameter ξ .

SPM thanks CSIR, Government of India, for his research fellowship.

-
- [1] Barone A and Paterno G 1982 *Physics and Applications of the Josephson Effect* (Wiley: New York).
 - [2] Golubov A A, Kupriyanov M Y and Il'ichev E 2004 Rev. Mod. Phys. **76** 411.
 - [3] Tsuei C C and Kirtley J R 2000 Rev. Mod. Phys. **72** 969.
 - [4] For review, Mackenzie A P and Maeno Y 2003 Rev. Mod. Phys. **75** 657.
 - [5] Bauer E, Hilscher G, Michor H, Paul C, Scheidt E W, Gribanov A, Seropegin Y, Noël H, Sigrist M, and Rogl P 2004 Phys. Rev. Lett. **92** 027003.
 - [6] Gor'kov L P and Rashba E I 2001 Phys. Rev. Lett. **87** 037004.
 - [7] Frigeri P A, Agterberg D F, Koga A, and Sigrist M 2004 Phys. Rev. Lett. **92** 097001.
 - [8] Rashba E I 1960 Sov. Phys. Solid State **2** 1109.
 - [9] Samokhin K V, Zijlstra E S, and Bose S K 2004 Phys. Rev. B **69** 094514.
 - [10] Sergienko I A and Curnoe S H 2004 Phys. Rev. B **70** 214510.
 - [11] Sumiyama A, Nakatsuji K, Tsuji Y, Oda Y, Yasuda T, Settai R and Onuki Y 2005 J. Phys. Soc. Japan **74** 3041.
 - [12] Ambegaokar V and Baratoff A 1963 Phys. Rev. Lett. **10** 486.
 - [13] Mahan G D 1990 *Many-Particle Physics* (Plenum Press: New York).

- [14] Duke C N 1969 *Tunneling in Solids*, Solid State Physics, Suppl. 10 (Academic: New York).
- [15] For example, $\delta \approx 0.4$ for $|T_{\mathbf{k}\mathbf{p}}|^2 \sim |T|^2 \hat{k}_\perp \hat{p}_\perp \Theta(\hat{k}_\perp \hat{p}_\perp)$ as suggested in Bruder C, Otterlo A van, and Zimanyi G T 1995 Phys. Rev. B **51** 12904, where Θ represents Heaviside function and subscript $\perp$ represents the direction of momentum along transverse to the junction.